

Compounded Anhydrous Sublingual Semaglutide Suspension

Please note if clinical success is not achieved by semaglutide alone, a prescriber or patient might consider the combination of semaglutide and tirzepatide as an alternative treatment.

Educational information can be found at www.subsema.com, both in written form as well as video.

Ingredients: Semaglutide and Submagna™ HMW

Warning: Semaglutide possesses a risk of thyroid c-cell tumors. This drug has been shown to cause thyroid cancer in some animals, however; it is unknown if Semaglutide causes thyroid c-cell tumors in humans. Semaglutide is contraindicated in patients with a personal or family history of multiple endocrine neoplasia syndrome type 2 (men 2) or thyroid cancer.

Uses: Semaglutide could be used to lower blood glucose in adults with type 2 diabetes. It could also be used for promoting weight loss in adjunct with diet and increased physical activity. It also is being studied in a number of other areas including, but not limited to, the following: type-1-diabetes, overindulgence, addiction, alcoholism, memory issues, kidney disease, liver disease, lung disease, nicotine intake and others. A massive number of clinical trials are underway related to semaglutide that can be found at www.clinicaltrials.gov.

How to Take this Medication: Use this medication as prescribed. Typical administration is to place the indicated volume under the patient's tongue and be allowed to absorb there for a minimum of 90 seconds then swallow – allowing to sit under the tongue for up to 5 minutes could even be more ideal. Take at least 30 minutes before the first food, drink, or medications of the day.

Side Effects: Most commonly reported side effects when taking Semaglutide (orally) are: nausea, abdominal pain, diarrhea, decreased appetite, vomiting, and constipation. Signs of an allergic reaction present as rash, hives, itching, blistered or peeling skin, tightness in chest, swelling of lips/tongue/throat, difficulty swallowing, or difficulty breathing. If you experience any of these symptoms, please contact your doctor immediately.

Precautions: Pancreatitis has been reported in patients taking semaglutide. Discontinue and contact your doctor if you experience severe stomach pain, back pain, and/or vomiting. Any signs of visual disturbances, low blood sugar, and changes in urine should be reported to your doctor before continuation of semaglutide.

Overdose: If an overdose is identified, please contact the pharmacy or call 911. Poison control can also be reached at 800.222.1222

Missed Dose: If a dose is missed, do not try to "make up" the dose. Go on to the next scheduled dose and move forward accordingly. Do not take 2 doses at the same time or extra doses.

Storage: Store this medication in a cool, dry place. avoid exposure to excessive heat, sunlight, or dampness.

Broken Arrow Location

811 W. New Orleans, Ste 101 • Broken Arrow, OK 74011



Directions:

From OK-51W / Broken Arrow Expressway:

- Take the Elm Pl exit, go 0.3 mi
- Turn south onto N Elm Pl, go 3.8 mi
- Turn right onto E 101st St/W New Orleans St, go 0.2 mi

From the Creek Turnpike:

- Turn the New Orleans St. exit, go 0.3 mi
- Turn right onto E 101st St/W New Orleans St, go 3.2 mi

Tulsa Location

4132 E. 51st St. • Tulsa, OK 74135



Directions:

From I-44:

- Take I-44 to exit 228 for E 51st St toward Harvard Ave, go 0.2 mi
- Turn left onto E 51st St, go 0.3 mi
- NextGenRx will be on your right

Vitality Concierge Membership

\$60 per year for each household



• **FREE SHIPPING**

• **FREE DELIVERY**

• **Concierge service**

• **Cost savings per prescription**

• **Unlimited hormone consultations**

• **Free antibiotics**
(limitations apply)

• **3 month supplies**

MyNextGenRx.com • Phone: 918.578.0031

Broken Arrow: 811 W. New Orleans, Ste 101, Broken Arrow, Ok 74011 | Fax: 918.524.9365 • Tulsa: 4132 E. 51st St., Tulsa, OK 74135 | Fax: 918.512.4255